

SEQUENCE LISTING

<110> KISIELOW, Malgorzata Anna
KLEINER, Sandra
NAGAMINE, Yoshikuni

<120> METHODS OF OBTAINING ISOFORM SPECIFIC
EXPRESSION IN MAMMALIAN CELLS

<130> 1-32330A/FMI

<150> GB 0201477.7

<151> 2002-01-23

<150> PCT/EP03/00611

<151> 2003-01-22

<160> 16

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 21

<212> RNA

<213> Artificial Sequence

<220>

<223> oligoribonucleotide pair used as siRNA; h/m-shc
siRNA from nt 677-697 (in the PTB domain)

<400> 1

cuacuugguu cgguacaugg g

21

<210> 2

<211> 22

<212> RNA

<213> Artificial Sequence

<220>

<223> oligoribonucleotide pair used as siRNA; h/m-shc
siRNA from nt 677-697 (in the PTB domain)

<400> 2

cauguaccga acccaaguag ga

22

<210> 3

<211> 21

<212> RNA

<213> Artificial Sequence

<220>

<223> oligonucleotide pair used for siRNA; h/m-shc
siRNA, p66-shc siRNA (from nt 236-256 in the CH2
domain)

<400> 3
 gaaugagucu cugucaucgu c 21

 <210> 4
 <211> 21
 <212> RNA
 <213> Artificial Sequence

 <220>
 <223> oligonucleotide pair used in siRNA; h/m-shc siRNA,
 p66-shc siRNA (from nt 236-256 in the CH2 domain)

 <400> 4
 cgaugacaga gacucauucc g 21

 <210> 5
 <211> 30
 <212> DNA
 <213> Mus musculus

 <220>
 <221> primer_bind
 <222> (1)...(30)
 <223> sense primer for mouse p46 ShcA cDNA

 <400> 5
 cggaattcat gggacctggg gtttctact 30

 <210> 6
 <211> 30
 <212> DNA
 <213> Mus musculus

 <220>
 <221> primer_bind
 <222> (1)...(30)
 <223> sense primer for mouse p52 ShcA cDNA

 <400> 6
 cggaattcat gaacaagctg agtggaggcg 30

 <210> 7
 <211> 36
 <212> DNA
 <213> Mus musculus

 <220>
 <221> primer_bind
 <222> (1)...(36)
 <223> sense primer for mouse p66 ShcA cDNA

 <400> 7
 cggaattcat ggatcttcta ccccccaagc cgaagt 36

 <210> 8
 <211> 31
 <212> DNA

<213> Mus musculus
 <220>
 <221> primer_bind
 <222> (1)...(31)
 <223> common antisense primer for ShcA cDNAs
 <400> 8
 cggaattcac actttccgat ccacgggttg c 31
 <210> 9
 <211> 40
 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> overlapping oligonucleotide pair for insertion
 into the KpnI-EcoRI site of pcDNA3
 <400> 9
 cccaccatgg cttaccata cgatgttcca gattacgctg 40
 <210> 10
 <211> 48
 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> overlapping oligonucleotide pair for insertion
 into the KpnI-EcoRI site of pcDNA3
 <400> 10
 aattcagcga attctggaac atcgtatggg taagccatgg tggggtac 48
 <210> 11
 <211> 27
 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> overlapping oligonucleotide pair used to mutate
 methionine 65 (start site for p52) to leucine in
 p66HA, resulting in p66HA-m1
 <400> 11
 ctcttcagg acctgaacaa gctgagt 27
 <210> 12
 <211> 28
 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> overlapping oligonucleotide pair used to mutate
 methionine 65 (start site for p52) to leucine in
 p66HA, resulting in p66HA-m1

<400> 12	
cactcagctt gttcagggtcc tggaggag	28
<210> 13	
<211> 27	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> overlapping oligonucleotide pair used to mutate the initiation sites for p46 in both p66HA-m1 and p52HA, resulting in p66HA-ML and p52HA-ML	
<400> 13	
ccaacgacaa agtcctggga cccgggg	27
<210> 14	
<211> 27	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> overlapping oligonucleotide pair used to mutate the initiation sites for p46 in both p66HA-m1 and p52HA, resulting in p66HA-ML and p52HA-ML	
<400> 14	
ccccgggtcc caggactttg tcgttgg	27
<210> 15	
<211> 34	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> overlapping oligonucleotide pair used to give p46HA-sm, p52HA-ML-sm and p66HA-ML-sm	
<400> 15	
ggggtttcct acttggtccg ctacatgggt tgtc	34
<210> 16	
<211> 34	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> overlapping oligonucleotide pair used to give p46HA-sm, p52HA-ML-sm and p66HA-ML-sm	
<400> 16	
cacaacccat gtagcggacc aagtaggaaa cccc	34